

TIER ONE ALGEBRA EXAM

- (1) Consider the matrix $A = \begin{pmatrix} -4 & 18 \\ -3 & 11 \end{pmatrix}$.
- (a) Find an invertible matrix P such that $P^{-1}AP$ is a diagonal matrix.
 - (b) Using the previous part of this problem, find a formula for A^n where A^n is the result of multiplying A by itself n times.
 - (c) Consider the sequences of numbers

$$a_0 = 1, b_0 = 0, a_{n+1} = -4a_n + 18b_n, b_{n+1} = -3a_n + 11b_n.$$

Use the previous parts of this problem to compute closed formulae for the numbers a_n and b_n .

- (2) Let P_2 be the vector space of polynomials with real coefficients and having degree less than or equal to 2. Define $D : P_2 \rightarrow P_2$ by $D(f) = f'$, that is, D is the linear transformation given by taking the derivative of the polynomial f . (You needn't verify that D is a linear transformation.)
- (3) Give an example of each of the following. (No justification required.)
- (a) A group G , a normal subgroup H of G , and a normal subgroup K of H such that K is not normal in G .
 - (b) A non-trivial perfect group. (Recall that a group is *perfect* if it has no non-trivial abelian quotient groups.)
 - (c) A field which is a three dimensional vector space over the field of rational numbers, \mathbb{Q} .
 - (d) A group with the property that the subset of elements of finite order is not a subgroup.
 - (e) A prime ideal of $\mathbb{Z} \times \mathbb{Z}$ which is not maximal.

- (4) Show that any field with four elements is isomorphic to

$$\frac{\mathbb{F}_2[t]}{(1 + t + t^2)}.$$

- (5) Let \mathbb{F}_p denote a field with p elements, p prime. Consider the ring

$$R = \frac{\mathbb{F}_p[x, y]}{\langle x^2 - 3, y^2 - 5 \rangle}$$

where $\langle x^2 - 3, y^2 - 5 \rangle$ denotes the ideal generated by $x^2 - 3$ and $y^2 - 5$. Show that R is not a field.

- (6) Let

$$R = \frac{\mathbb{C}[x, y]}{(x^2 + y^3)}$$

where $\mathbb{C}[x, y]$ is the polynomial ring over the complex numbers \mathbb{C} with indeterminates x and y . Similarly, let S be the subring of $\mathbb{C}[t]$ given by $\mathbb{C}[t^2, t^3]$.

- (a) Prove that R and S are isomorphic as rings.
 - (b) Let I be the ideal in R given by the residue classes of x and y . Prove that I is a prime ideal of R but not a principle ideal of R .
- (7) Suppose that $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a linear map.
- (a) Suppose $n = 2$ and $T^2 = -I$. Prove that T has no eigenvectors in \mathbb{R}^2 .
 - (b) Suppose $n = 2$ and $T^2 = I$. Prove that \mathbb{R}^2 has a basis consisting of eigenvectors of T .
 - (c) Suppose $n = 3$. Prove that T has an eigenvector in \mathbb{R}^3 . Give an example of an operator T such that T has an eigenvector in \mathbb{R}^3 , but \mathbb{R}^3 does not have a basis consisting of eigenvectors of T .
- (8) Let $p(x)$ and $q(x)$ be polynomials with rational coefficients such that $p(x)$ is irreducible over the field of rational numbers \mathbb{Q} . Let $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ be the complex roots of p , and suppose that $q(\alpha_1) = \alpha_2$. Prove that
- $$q(\alpha_i) \in \{\alpha_1, \alpha_2, \dots, \alpha_n\}$$
- for all $i \in \{2, 3, \dots, n\}$.
- (9) Let F be a field containing subfields F_{16} and F_{64} with 16 and 64 elements respectively. Find (with proof) the order of $F_{16} \cap F_{64}$.
- (10) Let G be a finite group and suppose H is a subgroup of G having index n . Show there is a normal subgroup K of G with $K \subset H$ and such that the order of K divides $n!$.
- (a) Find a matrix representing the linear function D in the basis $\{1, x, x^2\}$.
 - (b) Determine the eigenvalues and eigenvectors of D .
 - (c) Determine if P_2 has a basis such that D is represented by a diagonal matrix. Why or why not?
- (11) Suppose that W is a non-zero finite dimensional vector space over \mathbb{R} . Let T be a linear transformation of W to itself. Prove that there is a subspace U of W of dimension 1 or 2 such that $T(U) \subset U$ (i.e. U is an invariant subspace. Here $T(U)$ denotes the set $\{T(u) | u \in U\}$.)