

Tier 1 Analysis Exam
January 2000

1. Let Ω be an open set in \mathbb{R}^2 . Let u be a real-valued function on Ω . Suppose that for each point $a \in \Omega$ the partial derivatives $u_x(a)$ and $u_y(a)$ exist and are equal to zero.

(i) Prove that u is locally constant, *i.e.* for every point in Ω there is a neighborhood on which u is a constant function.

(ii) Prove that if Ω is connected, then u is a constant function on Ω .

2. Let S be the surface in the Euclidean space \mathbb{R}^3 given by the equation $x^2 + y^2 - z^2 = 1$, $0 \leq z \leq 1$, oriented so that the normal vector points away from the z -axis. Find

$\int_S \mathbf{F} \cdot d\mathbf{S}$, where \mathbf{F} is the vector field defined by

$$\mathbf{F}(x, y, z) = (-xy^2 + z^5, -x^2y, (x^2 + y^2)z) .$$

3. Let $f(x) = e^x - \cos x$ for $x \in \mathbb{R}$.

(i) Show that on a neighborhood around $x = 0$, f has an inverse function g with $g(0) = 0$.

(ii) Compute $g''(0)$.

(iii) Show that there exists $a > 0$ such that $f : (-a, \infty) \rightarrow (f(-a), \infty)$ is a homeomorphism.

4. For positive numbers k_1, k_2, k_3, \dots we define $[k_1] = \frac{1}{k_1}$, $[k_1, k_2] = \frac{1}{k_1 + [k_2]}$, $[k_1, k_2, k_3] = \frac{1}{k_1 + [k_2, k_3]}$, and inductively, $[k_1, \dots, k_{n+1}] = \frac{1}{k_1 + [k_2, \dots, k_{n+1}]}$. Prove that $\lim_{n \rightarrow \infty} [k_1, \dots, k_n]$ exists if $k_n \geq 2$ for all n .

5. Two circular holes of radius 1 *in* are drilled from the centers of two faces of a solid cube of volume 64 in^3 . Compute the volume of the remaining solid.

6. Let $\varphi_1, \varphi_2, \varphi_3, \dots$ be non-negative continuous functions on $[-1, 1]$ such that

(i) $\int_{-1}^1 \varphi_k(t) dt = 1$ for $k = 1, 2, 3, \dots$;

(ii) for every $\delta \in (0, 1)$ $\lim_{k \rightarrow \infty} \varphi_k = 0$ uniformly on $[-1, -\delta] \cup [\delta, 1]$.

Prove that for every continuous function $f : [-1, 1] \rightarrow \mathbb{R}$ we have

$$\lim_{k \rightarrow \infty} \int_{-1}^1 f(t) \varphi_k(t) dt = f(0) .$$

7. Suppose $\lim_{n \rightarrow \infty} a_n = a$, $\lim_{n \rightarrow \infty} b_n = b$, and let

$$c_n = \frac{a_1 b_n + a_2 b_{n-1} + \cdots + a_n b_1}{n} .$$

Prove that $\lim_{n \rightarrow \infty} c_n = ab$.

8. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a uniformly continuous function on \mathbb{R} . Prove that there exist positive constants A and B such that

$$|f(x)| \leq A|x| + B \quad \text{for all } x \in \mathbb{R} .$$

9. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function. Suppose $\lim_{x \rightarrow \infty} \frac{f(x)}{x} = 1$. Prove that there exists a sequence $\{x_n\}_{n=1}^{\infty}$ such that $\lim_{n \rightarrow \infty} x_n = \infty$ and $\lim_{n \rightarrow \infty} f'(x_n) = 1$.