

Tier I Analysis Exam

January 2009

Try to work all questions. They all are worth the same amount.

1. Assume f and g are uniformly continuous functions from $\mathbb{R}^1 \rightarrow \mathbb{R}^1$. If both f and g are also bounded, show that fg is also uniformly continuous. Then give an example to show that in general, if f and g are both uniformly continuous but not both bounded, then the product is not necessarily uniformly continuous. (Verify clearly that your counter-example is not uniformly continuous.)
2. Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are \mathcal{C}^2 functions, $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a \mathcal{C}^1 function and assume

$$f(0) = g(0) = 0, \quad f'(0) = g'(0) = h(0,0) = 1.$$

Show that the function $H : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$H(x, y) := \int_0^{f(x)} \int_0^{g(y)} h(s, t) ds dt + \frac{1}{2}x^2 + by^2$$

has a local minimum at the origin provided that $b > \frac{1}{2}$ while it has a saddle at the origin if $b < \frac{1}{2}$.

3. Let $H = \{(x, y, z) \mid z > 0 \text{ and } x^2 + y^2 + z^2 = R^2\}$, i.e. the upper hemisphere of the sphere of radius R centered at 0 in \mathbb{R}^3 . Let $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the vector field

$$F(x, y, z) = \left\{ x^2(y^2 - z^3), xzy^4 + e^{-x^2}y^4 + y, x^2y(y^2x^3 + 3)z + e^{-x^2-y^2} \right\}$$

Find $\int_H F \cdot \hat{n} dS$ where \hat{n} is the outward (upward) pointing unit surface normal and dS is the area element.

4. Let D be the square with vertices $(2,2)$, $(3,3)$, $(2,4)$, $(1,3)$. Calculate the improper integral

$$\int \int_D \ln(y^2 - x^2) dx dy .$$

5. Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}^1$ is a \mathcal{C}^4 function with the property that at some point $(x_0, y_0) \in \mathbb{R}^2$ all of the first and second order partial derivatives of f vanish. Suppose also that at least one partial derivative of third order does not vanish at (x_0, y_0) . Prove that f can have neither a local maximum nor a local minimum at this critical point.

6. Prove that the series $\sum_{n=1}^{\infty} \frac{nx}{1 + n^2 \log^2(n)x^2}$ converges uniformly on $[\varepsilon, \infty)$ for any $\varepsilon > 0$.

7. Suppose that $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is of class \mathcal{C}^1 , that $f(0, 0, 0) = 0$, and

$$f_2(0, 0, 0) \neq 0, \quad f_3(0, 0, 0) \neq 0, \quad \text{and} \quad f_2(0, 0, 0) + f_3(0, 0, 0) \neq -1$$

where $f_k = \frac{\partial f}{\partial x_k}$. Show that the system

$$f(x_1, f(x_1, x_2, x_3), x_3) = 0$$

$$f(x_1, x_2, f(x_1, x_2, x_3)) = 0$$

defines \mathcal{C}^1 functions $x_2 = \varphi(x_1)$, and $x_3 = \psi(x_1)$ for x_1 in a neighborhood of 0 satisfying

$$f(x_1, f(x_1, \varphi(x_1), \psi(x_1)), \psi(x_1)) = 0$$

$$f(x_1, \varphi(x_1), f(x_1, \varphi(x_1), \psi(x_1))) = 0.$$

8. For each $b \in [1, e]$, consider the sequence of real numbers governed by the recurrence relation

$$a_{n+1} = \left(\sqrt[n]{b}\right)^{a_n} \quad \text{for } n = 0, 1, 2, \dots \quad \text{with } a_0 = \sqrt[b]{b} \quad \text{i.e.} \quad \{\sqrt[b]{b}, \sqrt[n]{b}^{\sqrt[b]{b}}, \sqrt[n]{b}^{\sqrt[n]{b}^{\sqrt[b]{b}}}, \sqrt[n]{b}^{\sqrt[n]{b}^{\sqrt[n]{b}^{\sqrt[b]{b}}}}, \dots\}.$$

Show that this sequence converges and find the limit.

9. For each positive integer n , define $x_n : [-1, 1] \rightarrow \mathbb{R}$ by

$$x_n(t) = \begin{cases} -1 & \text{if } -1 \leq t \leq -1/n \\ nt & \text{if } -1/n < t < 1/n \\ 1 & \text{if } 1/n \leq t \leq 1 \end{cases}$$

(a) Show that $\{x_n\}$ is a Cauchy sequence in the metric space $(\mathcal{C}([-1, 1]), d)$, where $\mathcal{C}([-1, 1])$ denotes the set of continuous functions defined on $[-1, 1]$ and d denotes the metric given by

$$d(x, y) = \int_{-1}^1 |x(t) - y(t)| dt .$$

(b) Show that $(\mathcal{C}([-1, 1]), d)$ is not complete.