

**TIER I ANALYSIS EXAM**  
**AUGUST 2013**

Solve each of the following nine problems on a separate and clearly labeled sheet of paper. Fully justify your answers.

Notation:

- $\mathbb{R}$  is the set of real numbers
  - $\mathbb{R}^n$  is Euclidean space
  - $|x|$  is the Euclidean length of a vector  $x \in \mathbb{R}^n$ ; absolute value when  $n = 1$ .
- (1) Fix positive integers  $n, N$  and a bounded set  $A \subset \mathbb{R}^n$ . We use the notation

$$\overline{B}(a, r) = \{x \in \mathbb{R}^n : |x - a| \leq r\}, \quad a \in \mathbb{R}^n, r \geq 0.$$

Show that there exist  $a_1, a_2, \dots, a_N \in \mathbb{R}^n$  and numbers  $r_1, \dots, r_N \in [0, +\infty)$  such that

$$A \subset \bigcup_{k=1}^N \overline{B}(a_k, r_k)$$

and the sum  $\sum_{k=1}^N r_k^2$  is as small as possible. In other words, the set  $\{\sum_{k=1}^N r_k^2 : A \text{ can be covered with a collection } (\overline{B}(a_k, r_k))_{k=1}^N\}$  has a smallest element.

- (2) Is the sequence  $(\cos(\pi\sqrt{n^2 + n}))_{n=1}^{\infty}$  convergent?  
(3) For which values of  $x \in \mathbb{R}$  does the series

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{x + n}$$

converge? Is the convergence uniform on the interval  $(-1, 1)$ ?

- (4) Consider the functions  $f_n : [0, 1] \rightarrow \mathbb{R}$  defined by  $f_n(x) = (1 - x^n)^{2^n}$  for  $x \in [0, 1]$  and  $n \in \mathbb{N}$ . Prove that the limit  $\lim_{n \rightarrow \infty} f_n(x)$  exists for every  $x \in [0, 1]$ . Is the convergence uniform on  $[0, 1]$ ?  
(5) Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a Riemann integrable function and let  $\varepsilon > 0$ . Show that there exist continuous functions  $g, h : [0, 1] \rightarrow \mathbb{R}$  such that  $g(x) \leq f(x) \leq h(x)$  for all  $x \in [0, 1]$ , and

$$\int_0^1 (h(x) - g(x)) dx < \varepsilon.$$

Is the converse statement true?

- (6) Assume the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfies the property

$$f(x + t) \geq f(x) - t^2$$

for all real values of  $x$  and all positive values of  $t$ . Prove that  $f$  must be nondecreasing.

- (7) Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be everywhere differentiable, and assume that the Jacobian of  $f$  is not singular at any point  $x = (x_1, x_2) \in \mathbb{R}^2$ . Assume that  $|f(x)| \leq 1$  whenever  $|x| = 1$ , and prove that in fact  $|f(x)| \leq 1$  whenever  $|x| \leq 1$ .

(8) Compute

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{e^{-|x-y|^2}}{1+|x+y|^2} dx dy.$$

(9) Given a positive number  $r \neq 1$ , set  $C_r = \{(x, y) \in \mathbb{R}^2 : (x-1)^2 + y^2 = r\}$ . Calculate the line integral

$$\int_{C_r} \frac{x dy - y dx}{x^2 + y^2},$$

where  $C_r$  is oriented counterclockwise relative to  $(1, 0)$ .