

### Tier I Analysis January 2018

**Problem 1.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces. Let  $f : X \rightarrow Y$  be surjective such that

$$\frac{1}{2}d_X(x, y) \leq d_Y(f(x), f(y)) \leq 2d_X(x, y)$$

for all  $x, y \in X$ . Show that if  $(X, d_X)$  is complete, then also  $(Y, d_Y)$  is complete.

**Problem 2.** Show that

$$\lim_{n \rightarrow \infty} \left( 2\sqrt{n} - \sum_{k=1}^n \frac{1}{\sqrt{k}} \right)$$

exists.

**Problem 3.** Assume that *bitter* is a property of subsets of  $[0, 1]$  such that the union of two bitter sets is bitter. Subsets of  $[0, 1]$  that are not bitter are called *sweet*. Thus every subset of  $[0, 1]$  is either bitter or sweet. A *sweet spot* of a set  $A \subset [0, 1]$  is a point  $x_0 \in [0, 1]$  such that for every open set  $U \subset \mathbb{R}$  that contains  $x_0$ , the set  $A \cap U$  is sweet. Show that if  $A \subset [0, 1]$  is sweet, then  $A$  has a sweet spot.

**Problem 4.** Let  $f$  and  $g$  be periodic functions defined on  $\mathbb{R}$ , not necessarily with the same period. Suppose that

$$\lim_{x \rightarrow \infty} f(x) - g(x) = 0 .$$

Show that  $f(x) = g(x)$  for all  $x$ .

**Problem 5.** Let  $0 < x_n < 1$  be an infinite sequence of real numbers such that for all  $0 < r < 1$

$$\sum_{x_n < r} \log \frac{r}{x_n} \leq 1 .$$

Show that

$$\sum_{n=1}^{\infty} (1 - x_n) < \infty .$$

**Problem 6.** Suppose that the series  $\sum_{n=1}^{\infty} a_n$  converges conditionally. Show that the series

$$\sum_{n=3}^{\infty} n(\log n)(\log \log n)^2 a_n$$

diverges.

**Problem 7.** Find the absolute minimum of the function  $f(x, y, z) = xy + yz + zx$  on the set  $g(x, y, z) = x^2 + y^2 + z^2 = 12$ .

**Problem 8.** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a  $C^1$  map such that  $f^{-1}(y)$  is a finite set for all  $y \in \mathbb{R}^2$ . Show that the determinant  $\det df(x)$  of the Jacobi matrix of  $f$  cannot vanish on an open subset of  $\mathbb{R}^2$ .

**Problem 9.** A *regular surface* is given by a continuously differentiable map  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  so that the differential  $df_x : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  has rank 2 for all  $x \in \mathbb{R}^2$ . The *tangent plane*  $T_x$  is the 2-dimensional subspace  $df_x(\mathbb{R}^2) \subset \mathbb{R}^3$ . Assume that a vector field  $X$  in  $\mathbb{R}^3$  is orthogonal to  $T_x$  for all  $x$ , i.e.  $X(f(x)) \cdot Y = 0$  for all  $x \in \mathbb{R}^2$  and all  $Y \in T_x$ . Show that  $X \cdot (\nabla \times X) = 0$  at all points  $f(x)$ .

**Problem 10.** Let  $f(x, y)$  be a function defined on  $\mathbb{R}^2$  such that

- For any fixed  $x$ , the function  $y \mapsto f(x, y)$  is a polynomial in  $y$ ;
- For any fixed  $y$ , the function  $x \mapsto f(x, y)$  is a polynomial in  $x$ .

Show that  $f$  is a polynomial, i.e.

$$f(x, y) = \sum_{i,j=0}^N a_{ij} x^i y^j$$

with suitable  $a_{i,j} \in \mathbb{R}$ ,  $i, j = 0, \dots, N$ .