

TIER 1 ANALYSIS EXAM, JANUARY 2020

Write the solution to each of the following problems on a separate, clearly identified page. Each problem is graded on a scale of zero to ten.

Problem 1. Let $\{a_n\}_{n \in \mathbb{N}}$ be a sequence of nonnegative real numbers such that $\lim_{n \rightarrow \infty} a_n = 0$. Show that there exist infinitely many $n \in \mathbb{N}$ with the following property:

$$a_m \leq a_n \text{ for every } m \geq n.$$

Problem 2. Let $\{a_n\}_{n \in \mathbb{N}}$ be a sequence of nonnegative real numbers such that $\lim_{n \rightarrow \infty} a_n = 0$ and

$$|a_n - a_{n+1}| \leq \frac{1}{n^2} \text{ for every } n \in \mathbb{N}.$$

Prove that the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$ converges.

Problem 3. Denote by X the collection of all sequences $x = \{x_n\}_{n \in \mathbb{N}}$ with the property that $x_n \in [0, 1]$ for every $n \in \mathbb{N}$. Define a metric on X by

$$d(x, y) = \sup_{n \in \mathbb{N}} |x_n - y_n|, \quad x = \{x_n\}_{n \in \mathbb{N}}, y = \{y_n\}_{n \in \mathbb{N}} \in X.$$

Let $f : X \rightarrow \mathbb{R}$ be a uniformly continuous function. Show that f is bounded. (NOTE: Take for granted the fact that d is in fact a metric. The conclusion is not correct if f is just continuous.)

Problem 4. Define a sequence $\{a_n\}_{n \in \mathbb{N}}$ as follows:

$$a_1 = 1, a_2 = \sqrt{2}, a_3 = \sqrt{2\sqrt{3}}, \dots, a_n = \sqrt{2\sqrt{3\sqrt{\dots\sqrt{n}}}}, \quad n \geq 3.$$

Show that the sequence converges in \mathbb{R} .

Problem 5. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a differentiable function such that $f(0, 0) = 0$. Show that the improper integral

$$\iint_{x^2+y^2 \leq 1} \frac{f(x, y)}{(x^2 + y^2)^{4/3}} dx dy$$

converges, that is,

$$\lim_{\varepsilon \downarrow 0} \iint_{\varepsilon \leq x^2+y^2 \leq 1} \frac{f(x, y)}{(x^2 + y^2)^{4/3}} dx dy$$

exists.

Problem 6. Let $f : \mathbb{R} \rightarrow (0, +\infty)$ be a differentiable function such that $f'(x) > f(x)$ for every $x \in \mathbb{R}$.

(1) Show that there exists a constant $k > 0$ such that

$$(0.1) \quad \lim_{x \rightarrow \infty} f(x)e^{-kx} = +\infty.$$

(2) Find the least upper bound of the numbers k for which (0.1) can be proved.

Problem 7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that f' is uniformly continuous and $\lim_{x \rightarrow +\infty} f(x) = 2020$. Does the limit $\lim_{x \rightarrow +\infty} f'(x)$ necessarily exist? (NOTE: Prove if true, provide an example if false.)

Problem 8. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $f(x+1) = f(x)$ for every $x \in \mathbb{R}$. Define functions $f_n : \mathbb{R} \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, as follows:

$$f_1(x) = f(x), f_n(x) = \frac{1}{2}(f_{n-1}(x - 2^{-n}) + f_{n-1}(x + 2^{-n})), \quad x \in \mathbb{R}, n \geq 2.$$

Show that the sequence $\{f_n\}_{n \in \mathbb{N}}$ converges uniformly on \mathbb{R} .

Problem 9. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a continuously differentiable function. Suppose that the Jacobian determinant $\det Df(0,0)$ is equal to zero. Show that for every $\varepsilon > 0$ there exist $M, \delta > 0$ with the following property:

If B_r is the closed disk of radius $r < \delta$ centered at $(0,0)$, then $f(B_r)$ is contained in a rectangle with sides Mr and εr .